

## Review

# Effectiveness of Laser Application for Periodontal Surgical Therapy: Systematic Review and Meta-Analysis

Shabnam Behdin,\* Alberto Monje,\* Guo-Hao Lin,\* Bruce Edwards,\* Abdullah Othman,\* and Hom-Lay Wang\*

**Background:** Evidence has shown some improved clinical outcomes and morbidity reduction with the use of lasers for non-surgical periodontal therapy due to ablation, vaporization, hemostasis, and field sterilization. The purpose of this systematic review is to evaluate and compare studies involving lasers as monotherapy or adjunctive to surgical periodontal treatment.

**Methods:** Electronic and manual searches were conducted by two independent reviewers in several databases for articles written in English up to December 2014. Articles were included in this review if they reported outcomes of surgical periodontal therapy with and without the use of lasers. The primary outcome was probing depth (PD), and secondary outcomes were measured changes in clinical factors such as clinical attachment level (CAL) and gingival recession (GR). For the comparative studies included, the pooled weighted mean difference (WMD) and 95% confidence interval (CI) of each variable were calculated using random-effects meta-analysis.

**Results:** Eight and nine articles were included in the quantitative and qualitative analyses, respectively. Although low-to-moderate risk of bias was detected, high heterogeneity among studies was found. In flap surgery with or without laser treatment, there was no statistically significant difference in primary outcome. Similarly, in guided tissue regeneration (GTR)/enamel matrix derivative (EMD) with and without laser treatment, the WMD of PD was negligible; however, the GTR/EMD group showed better outcomes ( $P = 0.005$ ) than the laser group. Regarding the secondary outcomes, in the flap surgery group, the WMD of CAL gain was 1.34 mm, and the WMD of GR was  $-0.24$  mm; no significant difference was detected between groups. In GTR/EMD with and without laser treatment, the WMD of CAL gain was 0.10 mm and the WMD of recession was  $-0.18$  mm; again, no significant difference was detected between groups.

**Conclusions:** The available evidence is insufficient to support the effectiveness of dental lasers as an adjunct to resective or regenerative surgical periodontal therapy. However, precautions must be exercised when interpreting the results of this study because of the small sample size and high heterogeneity among studies. *J Periodontol 2015;86:1352-1363.*

### KEY WORDS

Evidence-based dentistry; lasers; meta-analysis; periodontal diseases; periodontitis; review.

\* Periodontics and Oral Medicine, School of Dentistry, University of Michigan, Ann Arbor, MI.

Periodontal disease, an inflammatory disease caused by opportunistic bacteria residing in the oral cavity, causes periodontal breakdown.<sup>1</sup> Nowadays, periodontal therapy involves not only arresting the disease process, but also regenerating the tissues lost during the disease process.<sup>1-5</sup> Intervention should be addressed with available treatment modalities. Non-surgical therapy (i.e., scaling and root planing [SRP])<sup>6</sup> remains an essential part of periodontal therapy. In SRP, diseased root surface debridement with different hand and ultrasonic instrumentation is performed to facilitate periodontal reattachment. Additionally, other non-surgical and surgical techniques such as subgingival curettage, gingivectomy, modified Widman flap, and apically positioned flap have been studied in an attempt to reduce probing depth (PD) and increase clinical attachment level (CAL).<sup>7-12</sup> Other treatment options aiming to regenerate tissues include guided tissue regeneration (GTR) or the application of growth factors, which have demonstrated varying degrees of success and predictability.<sup>1-5</sup> Regardless of the treatment modality, surgical periodontal therapy is often associated with pain and discomfort.

In the last decade, the use of lasers (light amplification by stimulated emission of radiation) has occupied part of the dialogue within periodontology and oral surgery because of several proposed advantages.<sup>6,13,14</sup> In the arena of periodontology, laser use as an adjunct to non-surgical therapy was demonstrated to enhance periodontal healing;<sup>15</sup> however, it is still a matter of debate.<sup>16</sup> Advantages over conventional periodontal flap surgeries include ablation, vaporization, hemostasis, pocket sterilization, and morbidity reduction.<sup>17,18</sup> Lasers are also advantageous in many aspects for periodontal treatment such as effective root surface debridement.<sup>17,18</sup> Lasers used in this arena include 1) CO<sub>2</sub>; 2) neodymium:yttrium-aluminum-garnet (Nd:YAG); 3) erbium:yttrium-aluminum-garnet (Er:YAG); and 4) diode lasers. The Er:YAG laser (ERL) has been reported to be the most promising for periodontal treatment. Its ability to ablate dental calculus effectively without overheating adjacent tissue has been demonstrated.<sup>17-20</sup> Some studies have even pointed out significant CAL gain in conjunction with non-surgical periodontal treatment.<sup>19-21</sup> Laser systems such as CO<sub>2</sub>, diode, and Nd:YAG have been used for oral soft tissue procedures (gingivectomy or frenectomy).<sup>22,23</sup> CO<sub>2</sub> lasers have been successfully used as an adjunctive tool to de-epithelialize the mucoperiosteal flap during traditional flap surgery.<sup>24</sup> Laser treatment, alone or in combination with mechanical treatment (SRP), has produced positive clinical outcomes with respect to CAL gain, decreased PD, and decreased bleeding on probing (BOP).<sup>20,25,26</sup> Similar findings were shown using ERL in the medium term for single-rooted teeth with chronic periodontitis (CP), and outcomes achieved

could be maintained over a 5-year period.<sup>27</sup> The pulsed Nd:YAG laser has been incorporated into protocols that attempt to regenerate lost alveolar bone, cementum, periodontal ligament (PDL), and connective tissue attachment around natural teeth.<sup>28-30</sup> A similar protocol is also used on compromised implant sites in an attempt to regenerate supporting bone.<sup>31,32</sup>

Although some evidence<sup>15</sup> has shown the benefits of using lasers for non-surgical periodontal therapy with regard to clinical outcomes, its effectiveness for surgical therapy remains unclear. Hence, the purpose of this systematic review is to evaluate and compare various lasers as monotherapy or adjunctive to surgical periodontal therapy.

## MATERIALS AND METHODS

### *Focused Question and Process*

The focused question for this review is, what is the effectiveness of laser adjunctive therapy in periodontal surgical treatment as measured by clinical parameters?

The patient, intervention, comparative, outcome (PICO)<sup>33</sup> process factors are as follows: P: dentate patients with periodontal disease (chronic or aggressive, moderate or severe), with or without bleeding, with or without exudate, with or without furcation involvement, and with or without gingival recession (GR); I: surgical periodontal treatment using a laser (CO<sub>2</sub>, Nd:YAG, Er:YAG, or diode) approved by the Food and Drug Administration for use in surgical (resective or regenerative) or non-surgical periodontal treatment with or without manual debridement; C: surgical resective or regenerative treatment using approaches other than laser, alone or with laser therapy; and O: benefits or lack thereof in PD reduction, CAL gain, radiographic bone (RB) gain, and decrease in GR, BOP, and exudate.


### *Eligibility Criteria*

Articles were included in this systematic review if they met the following inclusion criteria: prospective or retrospective, randomized or not, cohort or case series involving human patients in which outcomes of surgical periodontal therapy using lasers were compared with other surgical approaches. Several factors such as study design, number of patients included at the last follow-up assessment, number of defect sites, smoking or other systemic conditions that might alter the outcome, and type of procedure (including whether bone grafting material or barrier membrane was used) were extracted from the selected studies and analyzed. To address the aim of this study comprehensively, parameters such as PD, CAL, BOP, RB change, suppuration, and GR were further reported (Table 1).<sup>21,24,25,33-39</sup> These were subgrouped by initial and final values and change in percent or millimeters, depending on the parameter

**Table 1.**  
**RCT Studies Included in the Qualitative Assessment for Flap Surgery Versus Laser and Regenerative Surgery Versus Laser**

Treatment and Reference	Group		Patients (n)	Age (years)	Number of Defects		Type of Defect	Laser		Surgery Type		Follow-Up (months)
	Test	Control			Test	Control		Type	Wavelength	Test	Control	
OFD and laser Sculean et al. 2004 <sup>22</sup>	Flap + laser	Flap	23	ND	12	11	Deep IBD (PD >6 mm)	Er:YAG	160 mJ/pulse, 10 pulses/second	OFD	OFD	6
Gasparic and Skaleric 2007 <sup>27</sup>	Flap + laser	MWF	25	46.3 ± 9.2	73	73	Advanced CP (PD >6 mm)	Er:YAG	600 millisecond, 180 mJ/pulse at 20 Hz	MWF	MWF	60
Retzeppi et al. 2007 <sup>41</sup>	Laser doppler flowmetry in PPF	MWF	10	35 to 65	10	10	CP (PD ≥5 mm)	Laser doppler flowmetry		PPF	MWF	2
Crespi et al. 2011 <sup>42</sup>	CAF + laser	MWF	25	45	100	100	≥5 mm PD	CO <sub>2</sub>	10.6 μm	CAF + laser	MWF	180
Gokhale et al. 2012 <sup>37</sup>	Flap + laser	Flap	30	30 to 50	30	30	≥5 mm PD	CO <sub>2</sub>	980 nm, laser dose, 50 J/cm <sup>2</sup> /second	OFD	OFD	3
GTR and laser Schwarz et al. 2003 <sup>26</sup>	Laser + EMD CAF	EMD and CAF alone	22	ND	11	11	IBD	Er:YAG	160 mJ/pulse, 10 Hz	OFD + EMD EDTA + EMD	OFD + EDTA + EMD	6
Ozcelik et al. 2008 <sup>38</sup>	EMD + laser	EMD alone	22	31 to 49	22	22	IBD (PD ≥6 mm)	Diode	588 and 4 J/cm	OFD + EMD	OFD + EMD	12
Diliz et al. 2010 <sup>39</sup>	EMD + laser	EMD alone	21	33 to 46	21	21	IBD (PD ≥6 mm)	Nd:YAG	1 W, 10 Hz, 100 mJ, 1,064 nm	OFD + EMP	OFD + EMP	12
Doğan et al. 2014 <sup>40</sup>	GTR + laser	GTR alone	13	26 to 49	13	13	PD ≥5 mm	Nd:YAG	1,064 nm, 100 mW, 100 mJ	GTR	GTR	6

OFD = open flap debridement; RCT = randomized controlled trial; ND = not defined; IBD = infrabony defect; MWF = modified Widman flap; PPF = papilla preservation flap; CAF = coronally advanced flap; EMD = enamel matrix derivative.  
None of the study participants smoked or had other systemic conditions.



**Figure 1.**  
PRISMA flowchart for the screening process.

being studied. Excluded from the review were: 1) case reports or case series with <10 patients; 2) systematic reviews; 3) preclinical animal studies; 4) human trials studying the use of laser for non-surgical (i.e., SRP) therapy; and 5) human trials with missing information.

### Screening Process

Electronic and manual searches were conducted by two independent reviewers (SB and AM) in four databases (MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials, and Cochrane Oral Health Group Trials Register databases) for articles written in English up to December 2014. For the PubMed library, combinations of controlled terms (MeSH and Emtree) and key words were used whenever possible. The search terms used in title and abstract were as follows: (bone loss, periodontal[MeSH Terms] OR debridement, periodontal[MeSH Terms] OR debridement, periodontal epithelial[MeSH Terms] OR periodontal atrophy[MeSH Terms]) OR adult periodontitis[MeSH Terms] AND (periodontal[Other Term] OR periodontal flap[Other Term]) AND (carbon dioxide lasers[MeSH Terms] OR


erbium YAG laser[MeSH Terms] OR ablation, laser tissue[MeSH Terms] OR Doppler laser flowmetry[MeSH Terms] OR diode laser[MeSH Terms] OR lasers, neodymium doped yttrium aluminum garnet[MeSH Terms]) AND English language[filter] AND clinical trial[filter] AND humans[filter]. For the Embase and Cochrane Libraries, the terms were restricted to the aim of the study. The key terms applied for the Embase Library were as follows: laser AND periodontal AND surgery OR periodontal AND disease OR periodontal AND treatment AND clinical trials AND humans. For the Cochrane Library, laser periodontal surgery was applied in Title, Abstract, and Keywords in Trials. Additionally, to ensure a thorough screening process, a manual search was performed of periodontics-related journals, including *Journal of Dental Research*, *Journal of Clinical Periodontology*, *Journal of Periodontology*, and *The International Journal of Periodontics & Restorative Dentistry* from January 2014 to December 2014. References of included articles were screened to check all

available articles.

Two reviewers (AM and G-HL) designed and assessed the proposal for the project to make sure the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline was followed to minimize risk of bias in the screening process and provide a high level of evidence. PRISMA consists of a 27-item checklist and a four-phase flow diagram.<sup>34</sup>

### Qualitative Assessment

Criteria used to evaluate the quality of the selected randomized controlled trials (RCTs) were modified from the randomized clinical trial checklist of the Cochrane Center and the Consolidated Standards of Reporting Trials (CONSORT) statement,<sup>35</sup> which provides guidelines for the following parameters: 1) sequence generation; 2) allocation concealment method; 3) masking of the examiner; 4) address of incomplete outcome data; 5) free of selective outcome reporting; and 6) patient accounting at the end of the study. The degree of bias was categorized as low risk if all the criteria were met, moderate risk if one criterion was missing, and high



**Figure 2.** **A)** Meta-analysis for comparison of PD among selected studies in flap surgery with and without laser treatment. The overall WMD was 0.56 mm (95% CI = -0.56 to 1.67 mm, P = 0.33). **B)** Meta-analysis for comparison of PD among selected studies in GTR/EMD with and without laser treatment. The overall WMD was -0.01 mm (95% CI = -0.51 to 0.49 mm, P = 0.98). IV = independent variable.

risk if ≥2 criteria were missing.<sup>36</sup> Two independent reviewers (SB and AM) evaluated all the included articles.

**Statistical Analyses**

The primary outcome was PD reduction, and the secondary outcomes were CAL gain and GR reduction. BOP, suppuration, and RB change could not be meta-analyzed because of the high heterogeneity in studies reporting them.

Because of the variance in study designs and laser types, the analysis was divided into two subgroups: 1) flap surgery with and without laser treatment; and 2) GTR/enamel matrix derivative (EMD) with and without laser treatment. For the comparative studies included, the pooled weighted mean difference (WMD) and 95% confidence interval (CI) of each variable were calculated with random-effects meta-analysis by a computer program.<sup>†</sup> Forest plots were produced to graphically represent WMD and 95% CI in primary and secondary outcomes for the studies, using number of defect sites investigated as the analysis unit. In addition, heterogeneity among studies was assessed with the χ<sup>2</sup> test, with a P value < 0.05 representing significant heterogeneity. Funnel plots were also generated to report potential publication bias.

**RESULTS**

**Study Selection**

An initial screening yielded a total of 5,446 articles, of which 16 potentially relevant<sup>13,20,22,26-28,37-46</sup> articles were selected after an evaluation of titles and abstracts. The full texts of these articles were obtained and thoroughly evaluated. Nine articles<sup>22,26,27,37-42</sup> fulfilled the inclusion criteria and are included in the systematic review. Because of the lack of analytical data, one of these studies was excluded;<sup>41</sup> thus, only eight articles<sup>22,26,27,37-40,42</sup> are included in the quantitative analysis (Fig. 1). Details of all included studies were summarized in Table 1. Briefly, for open-flap debridement (OFD), 103 individuals were included, and for GTR/OFD + EMD, 78 individuals were included. Reasons for exclusion are displayed in supplementary Table 1 in online *Journal of Periodontology*.

**Study Quality**

All the studies included in the qualitative and quantitative analyses were RCTs. The RCT checklist of the Cochrane Center and the CONSORT statement were used to score studies' quality (see supplementary Table 2 in online *Journal of Periodontology*). Low

<sup>†</sup> Review Manager, v.5.0, The Nordic Cochrane Centre, Copenhagen, Denmark.

**Table 2.** Quantitative Assessment for Flap or Regenerative Surgery versus Laser: PD and CAL

Reference	Mean PD (mm)						Mean CAL (mm)					
	Initial		Final		Reduction		Initial		Final		Reduction	
	Test	Control	Test	Control	Test	Control	Test	Control	Test	Control	Test	Control
Scullean et al. 2004 <sup>22</sup>	7.8 ± 1.3	7.8 ± 0.8	4.1 ± 1.3	4.6 ± 1.6	3.7	3.2	9.8 ± 2.9	9.2 ± 1.2	7.2 ± 2.5	7.7 ± 1.6	ND	ND
Gaspirc and Skaleric 2007 <sup>27</sup>	5.63 ± 0.95	5.78 ± 0.82	2.84 ± 0.43	2.91 ± 0.55	2.79	2.87	6.14 ± 1.80	6.36 ± 2.00	3.97 ± 0.89	4.05 ± 0.89	ND	ND
Retzeppi et al. 2007 <sup>41</sup>	5.8 ± 0.1	5.7 ± 0.1	3.3 ± 0.1	2.8 ± 0.1	2.5 ± 0.2	2.8 ± 0.2	6.9 ± 0.2	6.7 ± 0.2	5.5 ± 0.2	5.4 ± 0.3	1.5 ± 2.1	1.3 ± 0.3
Crespi et al. 2011 <sup>42</sup>	7.91 ± 0.81	7.17 ± 0.33	4.00 ± 0.38	4.80 ± 0.45	ND	ND	8.71 ± 0.34	8.63 ± 0.22	3.61 ± 1.11	8.23 ± 0.63	ND	ND
Gokhale et al. 2012 <sup>37</sup>	6.03 ± 1.22	5.80 ± 1.19	2.97 ± 0.72	3.00 ± 0.95	ND	ND	11.07 ± 1.57	11.5 ± 1.97	9.70 ± 1.62	9.80 ± 1.77	ND	ND
Schwarz et al. 2003 <sup>26</sup>	8.6 ± 1.2	8.1 ± 0.8	4.6 ± 0.8	4.0 ± 0.5	ND	ND	10.7 ± 1.3	10.4 ± 1.1	7.5 ± 1.4	7.1 ± 1.2	3.2	3.3
Ozcelik et al. 2008 <sup>38</sup>	6.6 (5.9 ± 6.8)	6.5 (5.9 ± 6.7)	1.3 (1.2 ± 2.3)	1.7 (1.3 ± 3.3)	ND	ND	7.2 (6.8 ± 7.5)	7.2 (6.5 ± 7.5)	3.1 (2.4 ± 3.7)	3.2 (2.8 ± 4.7)	ND	ND
Dilsiz et al. 2010 <sup>39</sup>	7.3 ± 0.6	7.3 ± 0.7	3.3 ± 0.4	3 ± 0.4	ND	ND	9.5 ± 0.7	9.3 ± 0.8	6.9 ± 0.7	6.4 ± 0.5	ND	ND
Doğan et al. 2014 <sup>40</sup>	6.01 ± 0.47	5.95 ± 0.43	2.63 ± 0.29	2.93 ± 1.98	3.38 ± 0.27	3.02 ± 0.42	7.35 ± 0.48	7.43 ± 0.60	4.46 ± 0.57	5.18 ± 0.46	2.91 ± 0.43	2.26 ± 0.34

ND = not defined.

(55.5% of studies) to moderate (44.5% of studies) estimated potential risk of bias was found from the studies included in the qualitative appraisal.

**Results of the Primary Outcome (PD)**

In flap surgery with and without laser treatment, the WMD of PD was 0.56 mm (95% CI = -0.56 to 1.67 mm, *P* = 0.33) (Fig. 2A). No significant difference was detected among groups. A high degree of heterogeneity (*P* < 0.001 for  $\chi^2$  test) was noted.

In GTR/EMD with and without laser treatment, the WMD of PD was -0.01 mm (95% CI = -0.51 to 0.49 mm, *P* = 0.98) (Fig. 2B). No significant difference was detected between groups. A high degree of heterogeneity (*P* < 0.001 for  $\chi^2$  test) was noted. Interestingly, in the EMD subgroup, the meta-analysis showed a significantly favorable outcome toward the EMD/GTR-only group (*P* = 0.005), with a low degree of heterogeneity between studies (*P* = 0.56 for  $\chi^2$  test).


Data for the initial, final, and reduction in mean PD for the included studies are presented in Table 2.

**Results of the Secondary Outcome (CAL gain and GR decrease)**

In flap surgery with and without laser treatment, the WMD of CAL gain was 1.34 mm (95% CI = -2.02 to 4.70 mm, *P* = 0.44) (Fig. 3A). No significant difference was detected among groups. The *P* value for  $\chi^2$  test was < 0.001, indicating a high heterogeneity among studies. In GTR/EMD with and without laser treatment, the WMD of CAL gain was 0.10 mm (95% CI = -0.64 to 0.85 mm, *P* = 0.78) (Fig. 3B). No significant difference was detected among groups. The *P* value for  $\chi^2$  test was < 0.001, indicating a high heterogeneity among studies. However, in the EMD subgroup, the meta-analysis showed a significantly favorable outcome toward EMD/GTR without the use of laser (*P* = 0.03), with a low degree of heterogeneity among studies (*P* value for  $\chi^2$  test 0.69). Data for the initial, final, and reduction in mean CAL for the included studies are presented in Table 2.

The WMD of GR was -0.24 mm (95% CI = -0.82 to 0.35 mm, *P* = 0.42) (Fig. 4A). No significant difference was detected among groups. The *P* value for  $\chi^2$  test was 0.18, indicating a moderate degree of heterogeneity among studies. Additionally, the WMD of GR was -0.18 mm (95% CI -0.42 to 0.07 mm, *P* = 0.15) in the comparison of EMD/GTR with and without the use of lasers (Fig. 4B). No significant difference was detected. The *P* value for  $\chi^2$  test was 0.59, indicating a low degree of heterogeneity among studies. Data for initial, final, and reduction in mean BOP, mean GR, and discomfort are presented in Table 3.

Funnel plots evaluating the publication bias of each parameter are shown in supplementary Figures 1 through 6 in online *Journal of Periodontology*.



### Figure 3.

**A)** Meta-analysis for comparison of CAL gain among selected studies in flap surgery with and without laser treatment. The overall WMD was 1.34 mm (95% CI = -2.02 to 4.70 mm,  $P = 0.44$ ). **B)** Meta-analysis for comparison of CAL gain among selected studies in GTR/EMD with and without laser treatment. The overall WMD was 0.10 mm (95% CI -0.64 to 0.85 mm,  $P = 0.78$ ). IV = independent variable.

## DISCUSSION

The effective applicability of lasers in the treatment of periodontal disease according to their theoretic advantages (i.e., ablation or vaporization, hemostasis, and sterilization effect) compared to conventional therapy is currently a very controversial topic in clinical periodontics. Findings from this study show that for regeneration and PD reduction surgical procedures, adjunctive use of lasers offers no significant clinical advantages in CAL gain and PD reduction compared with conventional approaches. These results must be cautiously interpreted, since limitations exist in regard to sample size and heterogeneity among studies.


### Summary of Main Findings

Different types of laser, including CO<sub>2</sub>, Nd:YAG, Er:YAG, and diode, have been used for periodontal disease treatment and hard and soft tissue management.<sup>15,16,47</sup> Laser irradiation, at low level, stimulates surrounding tissue cells and results in reduction of inflammation, higher tissue regeneration, better tissue attachment, and even increased lymph flow, as well as less postoperative pain, once the scattered beams penetrate into pockets.<sup>48</sup> Smooth and flat root surfaces with sealed dentinal tubules as well as bacterial elimination can be obtained using CO<sub>2</sub> lasers (in

defocused pulsed mode with power of 2W), which in turn enhance fibroblast attachment.<sup>49</sup> A diode laser with wavelength of 810 or 910 to 980 nm is a useful setup for soft tissue management (coagulating and cutting gingiva or oral mucosa, sulcular debridement) and has antibacterial effects.<sup>50</sup> It must be noted that CO<sub>2</sub>, Nd:YAG, and diode lasers are not effective in treating or dealing with hard tissue surfaces in terms of calculus removal.<sup>23,49</sup> In contrast, the ERL has the ability to effectively ablate dental calculus without creating thermal damage to adjacent tissue.<sup>19</sup> ERL has a wavelength of 2,940 nm, close to the absorption coefficient of water, making it safe and free of thermal side effects.<sup>23</sup> Moreover, ERL has antimicrobial effects and works to remove lipopolysaccharide from root surfaces.<sup>23</sup> In the present study, it was not the aim to compare the effectiveness of laser subtypes, and no particular one was found to be superior.

### Agreements and Disagreements With Previous Studies

In contrast to these findings, a long-term follow-up study<sup>42</sup> showed that in deep pockets (PD  $\geq 7$  mm), a coronally advanced flap combined with CO<sub>2</sub> laser provides greater PD reduction compared with a modified Widman flap procedure. A similar trend was also noted in CAL assessment. It was then concluded



**Figure 4.** **A)** Meta-analysis for comparison of GR among selected studies in flap surgery with and without laser treatment. The overall WMD was  $-0.24$  mm (95% CI =  $-0.82$  to  $0.35$  mm,  $P = 0.42$ ). **B)** Meta-analysis for comparison of GR among selected studies in GTR/EMD with and without laser treatment. The overall WMD was  $-0.18$  mm (95% CI =  $-0.42$  to  $0.07$  mm,  $P = 0.15$ ). IV = independent variable.

that CO<sub>2</sub> lasers might provide meaningful clinical benefits in the treatment of deep pockets. These findings have been partially attributed to the ability of the CO<sub>2</sub> laser to remove the smear layer and eliminate bacterial cells from diseased root surfaces.<sup>51</sup> Also, myofibroblasts, which are suggested to be responsible for tissue contraction, are less abundant after treatment with CO<sub>2</sub> lasers,<sup>51</sup> which may be another rationale for better CAL gain in the test group. Nonetheless, the findings may be attributed not just to the laser, but also to the study design, in which no similar surgical procedures were evaluated. Moreover, this specific study showed no crater formation or root surface damage resulting from laser therapy. These results were in contrast to previously reported studies.<sup>51-54</sup>

Other studies<sup>26,37,39</sup> are in agreement with the present findings. In a comparative evaluation<sup>37</sup> of the efficiency of the diode laser as an adjunct to mechanical debridement versus conventional mechanical debridement in periodontal open flap surgery, no difference was found between laser-treated and non-laser-treated groups with regard to clinical parameters. However, it was reported<sup>37</sup> that the use of laser treatment adjunctive to open flap debridement provides a beneficial effect by reduction of anaerobic bacterial colonies within the sulci. The antiseptic potential of

lasers was explained based on laser energy disrupting the protective mechanism of the organisms.<sup>55</sup> The visual analog scale showed that patients tolerated application of the diode laser; nonetheless, its application did not make a difference in pain perception.<sup>55</sup> Similar findings were reported by another group.<sup>37</sup>

Dental lasers have slight advantages but also some inherent drawbacks that have to be pointed out, including possible damage to bone owing to overheating and ineffectiveness in removing calculus. Hence, intermittent laser application and correct laser settings are key factors to reduce potential damage. Another investigation using ERL for the surgical treatment of chronic periodontal disease<sup>26</sup> found no statistically significant clinical differences compared with resective procedures.

Dental lasers have been applied and investigated for regenerative procedures (i.e., GTR).<sup>38,56</sup> In a short-term study,<sup>56</sup> patients with CP who had intrabony defects and were receiving treatment with GTR in combination with low-level laser therapy demonstrated statistically better improvements in PD reduction, CAL gain, and decreased GR at 6 months versus the group who received GTR alone. Additionally, a significant reduction of the sulcular bleeding index favored the laser-treated group. These findings



**Table 3.** Quantitative Assessment for Flap or Regenerative Surgery versus Laser: BOP, Recession, and Discomfort

Reference	Mean BOP (%)				Mean Recession (mm)								Discomfort of Procedure
	Initial		Final		Initial		Final		Increase		Control	Test	
	Test	Control	Test	Control	Test	Control	Test	Control	Test	Control			
Sculean et al. 2004, <sup>22</sup>	40	44	15	18	25	26	2.2 ± 1.6	1.5 ± 1.0	3.1 ± 2.0	3.2 ± 1.3	0.9	1.7	ND
Gasparic and Skaleric 2007 <sup>27</sup>	40.2	39.4	15.6	16.7	15.6	22.7	0.61 ± 0.25	0.58 ± 0.30	1.2 ± 0.39	1.25 ± 0.45	ND	ND	ND
Retzepi et al. 2007 <sup>41</sup>	ND	ND	ND	ND	ND	ND	1.1 ± 0.1	1.1 ± 0.1	2.2 ± 0.1	2.6 ± 0.1	-1.1 ± 0.1	-1.5 ± 0.2	ND
Crespi et al. 2011 <sup>42</sup>	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	Similar in both groups
Gokhale et al. 2012 <sup>37</sup>	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Schwarz et al. 2003 <sup>26</sup>	65	57	30	31	35	26	2.1 ± 0.9	2.3 ± 0.9	2.9 ± 1.0	3.1 ± 1.0	-0.8	-0.8	ND
Ozcelik et al. 2008 <sup>38</sup>	ND	ND	ND	ND	ND	ND	1.9 (1.1 ± 2.4)	2.3 (1.8 ± 3.0)	2.3 (1.1 ± 2.8)	2.9 (2.4 ± 3.8)	ND	ND	ND
Dilisz et al. 2010 <sup>39</sup>	100	100	9.52	4.76	90.48	95.24	2.2 ± 0.5	2 ± 0.4	3.6 ± 0.5	3.4 ± 0.3	ND	ND	ND
Doğan et al. 2014 <sup>40</sup>	ND	ND	ND	ND	ND	ND	1.33 ± 0.42	1.48 ± 0.55	1.82 ± 0.52	2.24 ± 0.43	ND	ND	ND

ND = not defined.

were partially corroborated with the same laser type by other studies carrying out other periodontal regenerative approaches.<sup>38,56</sup> Thus, it seems that low-level laser therapy as an adjunct to GTR may improve short-term clinical outcomes. Some in vitro studies were conducted to evaluate the responsiveness of PDL cells to low-level laser therapy.<sup>57-59</sup> Less collagen breakdown after laser irradiation was described in terms of plasminogen activator plasmin proteolytic system inhibition with the use of lasers.<sup>60</sup> Additionally, sites treated with an 809-nm laser showed higher fibroblast proliferation as well as increased production of basic fibroblast growth factor.<sup>59</sup> This can be mainly attributed to the stimulatory effect of laser therapy on PDL cell regeneration.<sup>59</sup>

Contrary to previous findings<sup>38,40,42</sup> and in agreement with ours, Dilsiz et al.<sup>39</sup> compared the clinical outcomes of enamel matrix proteins (EMP) alone and combined with Nd:YAG laser in the treatment of infrabony defects. They reported that the use of Nd:YAG laser did not have superiority over EDTA as a root conditioning agent.<sup>39</sup> Along these lines, it is important to mention that consensus is lacking that EMP as a biologic agent enhances clinical outcomes compared to GTR procedures using barrier membranes.<sup>61-64</sup> Also noteworthy is that there are still controversies in the literature regarding the effectiveness of root conditioning agents.<sup>65-67</sup> Other studies demonstrated the efficacy of laser application as a root biomodification instrument to achieve better periodontal regenerative outcome by means of new attachment formation.<sup>68-70</sup> Cementum biocompatibility alteration using Nd:YAG laser for more predictable new attachment after periodontium regeneration has also been demonstrated.<sup>70</sup>

**Limitations of the Study**

Finally, it is important to highlight some evident limitations found in the qualitative assessment of articles using dental lasers as an adjunct to regeneration. For instance, defect morphology can potentially affect the final outcome. Steffensen and Webert showed statistically better results for defects angled <45 degrees. As a matter of fact, wider defects had higher RB loss.<sup>71</sup> Likewise, Cortellini and Tonetti demonstrated

that in scenarios of radiographic angulation  $\leq 25$  degrees, presence of a deep infrabony defect ( $\geq 3$  mm), and  $\geq 1$  mm of gingival thickness over the site, the most consistent CAL and bone gains were achieved, regardless of the number of bony walls.<sup>72</sup> Many factors contribute to the results of any regenerative procedure, and precautions must be exercised when interpreting the present findings.

Moreover, it is essential to understand the limitations of the present review based on the heterogeneity of the studies included. Certainly, all the included studies were RCTs; nonetheless, as evaluated by the CONSORT statement, 44.5% have moderate risk of bias, mainly on two sections: 1) masking examiner and 2) selective outcome reporting. In such a controversial topic, where conflict of interest might play a major role in the outcome, studies on this matter must be cautiously interpreted to avoid misleading readers.

## CONCLUSIONS

Within the limits of this study, this systematic review fails to support the effectiveness of dental lasers as an adjunct to resective or regenerative surgical periodontal therapy, owing to the lack of significant superiority over conventional approaches. Precautions must be exercised when interpreting the results of this study because of the small sample size and high heterogeneity among studies. Therefore, more randomized clinical trials examining different laser types and wavelengths are required to obtain stronger conclusions in this regard.

## ACKNOWLEDGMENTS

This paper was partially supported by the University of Michigan Periodontal Graduate Student Research Fund. H-LW receives financial support for lecturing from Morita Corp. (Osaka, Japan). The other authors do not have any financial interests, either directly or indirectly, in the products or information listed in the paper.

## REFERENCES

- Bowers GM, Granet M, Stevens M, et al. Histologic evaluation of new attachment in humans. A preliminary report. *J Periodontol* 1985;56:381-396.
- Mellonig JT. Decalcified freeze-dried bone allograft as an implant material in human periodontal defects. *Int J Periodontics Restorative Dent* 1984;4:40-55.
- Sanders JJ, Sepe WW, Bowers GM, et al. Clinical evaluation of freeze-dried bone allografts in periodontal osseous defects. Part III. Composite freeze-dried bone allografts with and without autogenous bone grafts. *J Periodontol* 1983;54:1-8.
- Quintero G, Mellonig JT, Gambill VM, Pelleu GB Jr. A six-month clinical evaluation of decalcified freeze-dried bone allografts in periodontal osseous defects. *J Periodontol* 1982;53:726-730.
- Bowers GM, Schallhorn RG, Mellonig JT. Histologic evaluation of new attachment in human intrabony defects. A literature review. *J Periodontol* 1982;53:509-514.
- Yukna RA. Lasers in periodontal therapy. *Today's FDA* 2011;23:40-41.
- Ramfjord SP, Nissle RR. The modified Widman flap. *J Periodontol* 1974;45:601-607.
- Ramfjord SP. Present status of the modified Widman flap procedure. *J Periodontol* 1977;48:558-565.
- Ramfjord SP, Caffesse RG, Morrison EC, et al. 4 modalities of periodontal treatment compared over 5 years. *J Clin Periodontol* 1987;14:445-452.
- Carnevale G, Kaldahl WB. Osseous resective surgery. *Periodontol 2000* 2000;22:59-87.
- Becker W, Becker BE, Caffesse R, et al. A longitudinal study comparing scaling, osseous surgery, and modified Widman procedures: Results after 5 years. *J Periodontol* 2001;72:1675-1684.
- Kaldahl WB, Kalkwarf KL, Patil KD, Dyer JK, Bates RE Jr. Evaluation of four modalities of periodontal therapy. Mean probing depth, probing attachment level and recession changes. *J Periodontol* 1988;59:783-793.
- Nevins ML, Camelo M, Schupbach P, Kim SW, Kim DM, Nevins M. Human clinical and histologic evaluation of laser-assisted new attachment procedure. *Int J Periodontics Restorative Dent* 2012;32:497-507.
- Nevins M, Nevins ML, Yamamoto A, et al. Use of Er:YAG laser to decontaminate infected dental implant surface in preparation for reestablishment of bone-to-implant contact. *Int J Periodontics Restorative Dent* 2014;34:461-466.
- Schwarz F, Aoki A, Becker J, Sculean A. Laser application in non-surgical periodontal therapy: A systematic review. *J Clin Periodontol* 2008;35(Suppl. 8):29-44.
- Sgolastra F, Petrucci A, Gatto R, Monaco A. Efficacy of Er:YAG laser in the treatment of chronic periodontitis: Systematic review and meta-analysis. *Lasers Med Sci* 2012;27:661-673.
- Aoki A, Ando Y, Watanabe H, Ishikawa I. In vitro studies on laser scaling of subgingival calculus with an erbium:YAG laser. *J Periodontol* 1994;65:1097-1106.
- Israel M, Cobb CM, Rossmann JA, Spencer P. The effects of CO<sub>2</sub>, Nd:YAG and Er:YAG lasers with and without surface coolant on tooth root surfaces. An in vitro study. *J Clin Periodontol* 1997;24:595-602.
- Schwarz F, Pütz N, Georg T, Reich E. Effect of an Er:YAG laser on periodontally involved root surfaces: An in vivo and in vitro SEM comparison. *Lasers Surg Med* 2001;29:328-335.
- Schwarz F, Sculean A, Georg T, Reich E. Periodontal treatment with an Er:YAG laser compared to scaling and root planing. A controlled clinical study. *J Periodontol* 2001;72:361-367.
- Watanabe H, Ishikawa I, Suzuki M, Hasegawa K. Clinical assessments of the erbium:YAG laser for soft tissue surgery and scaling. *J Clin Laser Med Surg* 1996;14:67-75.
- Sculean A, Schwarz F, Berakdar M, Windisch P, Arweiler NB, Romanos GE. Healing of intrabony defects following surgical treatment with or without an Er:YAG laser. *J Clin Periodontol* 2004;31:604-608.
- Sculean A, Schwarz F, Berakdar M, Romanos GE, Arweiler NB, Becker J. Periodontal treatment with an Er:YAG laser compared to ultrasonic instrumentation: A pilot study. *J Periodontol* 2004;75:966-973.
- Centy IG, Blank LW, Levy BA, Romberg E, Barnes DM. Carbon dioxide laser for de-epithelialization of periodontal flaps. *J Periodontol* 1997;68:763-769.

25. Tomasi C, Schander K, Dahlén G, Wennström JL. Short-term clinical and microbiologic effects of pocket debridement with an Er:YAG laser during periodontal maintenance. *J Periodontol* 2006;77:111-118.
26. Schwarz F, Sculean A, Georg T, Becker J. Clinical evaluation of the Er:YAG laser in combination with an enamel matrix protein derivative for the treatment of intrabony periodontal defects: A pilot study. *J Clin Periodontol* 2003;30:975-981.
27. Gaspirc B, Skaleric U. Clinical evaluation of periodontal surgical treatment with an Er:YAG laser: 5-year results. *J Periodontol* 2007;78:1864-1871.
28. Yukna RA, Carr RL, Evans GH. Histologic evaluation of an Nd:YAG laser-assisted new attachment procedure in humans. *Int J Periodontics Restorative Dent* 2007;27:577-587.
29. Fornaini C, Rocca JP, Bertrand MF, Merigo E, Nammour S, Vescovi P. Nd:YAG and diode laser in the surgical management of soft tissues related to orthodontic treatment. *Photomed Laser Surg* 2007;25:381-392.
30. de Andrade AK, Feist IS, Pannuti CM, Cai S, Zzell DM, De Micheli G. Nd:YAG laser clinical assisted in class II furcation treatment. *Lasers Med Sci* 2008;23:341-347.
31. Deppe H, Horch HH. Laser applications in oral surgery and implant dentistry. *Lasers Med Sci* 2007;22:217-221.
32. Giannini R, Vassalli M, Chellini F, Polidori L, Dei R, Giannelli M. Neodymium:yttrium aluminum garnet laser irradiation with low pulse energy: A potential tool for the treatment of peri-implant disease. *Clin Oral Implants Res* 2006;17:638-643.
33. Stone PW. Popping the (PICO) question in research and evidence-based practice. *Appl Nurs Res* 2002;15:197-198.
34. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med* 2009;6:e1000100.
35. Hopewell S, Clarke M, Moher D, et al; CONSORT Group. CONSORT for reporting randomized controlled trials in journal and conference abstracts: Explanation and elaboration. *PLoS Med* 2008;5:e20.
36. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: Updated guidelines for reporting parallel group randomised trials. *J Pharmacol Pharmacother* 2010;1:100-107.
37. Gokhale SR, Padhye AM, Byakod G, Jain SA, Padbidri V, Shivaswamy S. A comparative evaluation of the efficacy of diode laser as an adjunct to mechanical debridement versus conventional mechanical debridement in periodontal flap surgery: A clinical and microbiological study. *Photomed Laser Surg* 2012;30:598-603.
38. Ozelik O, Cenk Haytac M, Seydaoglu G. Enamel matrix derivative and low-level laser therapy in the treatment of intra-bony defects: A randomized placebo-controlled clinical trial. *J Clin Periodontol* 2008;35:147-156.
39. Dilsiz A, Canakci V, Aydin T. The combined use of Nd:YAG laser and enamel matrix proteins in the treatment of periodontal intrabony defects. *J Periodontol* 2010;81:1411-1418.
40. Doğan GE, Demir T, Orbak R. Effect of low-level laser on guided tissue regeneration performed with equine bone and membrane in the treatment of intrabony defects: A clinical study. *Photomed Laser Surg* 2014;32:226-231.
41. Retzepi M, Tonetti M, Donos N. Comparison of gingival blood flow during healing of simplified papilla preservation and modified Widman flap surgery: A clinical trial using laser Doppler flowmetry. *J Clin Periodontol* 2007;34:903-911.
42. Crespi R, Cappare P, Gherlone E, Romanos GE. Comparison of modified Widman and coronally advanced flap surgery combined with CO<sub>2</sub> laser root irradiation in periodontal therapy: A 15-year follow-up. *Int J Periodontics Restorative Dent* 2011;31:641-651.
43. White JM, Goodis HE, Rose CL. Use of the pulsed Nd:YAG laser for intraoral soft tissue surgery. *Lasers Surg Med* 1991;11:455-461.
44. Rotundo R, Nieri M, Cairo F, et al. Lack of adjunctive benefit of Er:YAG laser in non-surgical periodontal treatment: A randomized split-mouth clinical trial. *J Clin Periodontol* 2010;37:526-533.
45. Belal MH, Watanabe H. Comparative study on morphologic changes and cell attachment of periodontitis-affected root surfaces following conditioning with CO<sub>2</sub> and Er:YAG laser irradiations. *Photomed Laser Surg* 2014;32:553-560.
46. Miyazaki A, Yamaguchi T, Nishikata J, et al. Effects of Nd:YAG and CO<sub>2</sub> laser treatment and ultrasonic scaling on periodontal pockets of chronic periodontitis patients. *J Periodontol* 2003;74:175-180.
47. Sgolastra F, Petrucci A, Gatto R, Monaco A. Effectiveness of laser in dentinal hypersensitivity treatment: A systematic review. *J Endod* 2011;37:297-303.
48. Gokhale R, Agarkar S, Debgupta J, et al. Laser synthesized super-hydrophobic conducting carbon with broccolitype morphology as a counter-electrode for dye sensitized solar cells. *Nanoscale* 2012;4:6730-6734.
49. Crespi R, Barone A, Covani U, Ciaglia RN, Romanos GE. Effects of CO<sub>2</sub> laser treatment on fibroblast attachment to root surfaces. A scanning electron microscopy analysis. *J Periodontol* 2002;73:1308-1312.
50. Sanz-Moliner JD, Nart J, Cohen RE, Ciancio SG. The effect of an 810-nm diode laser on postoperative pain and tissue response after modified Widman flap surgery: A pilot study in humans. *J Periodontol* 2013;84:152-158.
51. Tucker D, Cobb CM, Rapley JW, Killoy WJ. Morphologic changes following in vitro CO<sub>2</sub> laser treatment of calculus-laden root surfaces. *Lasers Surg Med* 1996;18:150-156.
52. González M, Banderas JA, Rodríguez V, Castaño VM. Particle-induced X-ray emission and scanning electron microscopic analyses of the effects of CO<sub>2</sub> laser irradiation on dentinal structure. *J Dent* 1999;27:595-600.
53. Sasaki KM, Aoki A, Masuno H, Ichinose S, Yamada S, Ishikawa I. Compositional analysis of root cementum and dentin after Er:YAG laser irradiation compared with CO<sub>2</sub> laser and intact roots using Fourier transformed infrared spectroscopy. *J Periodontol Res* 2002;37:50-59.
54. Lan WH, Chen KW, Jeng JH, Lin CP, Lin SK. A comparison of the morphological changes after Nd:YAG and CO<sub>2</sub> laser irradiation of dentin surfaces. *J Endod* 2000;26:450-453.
55. Read RP, Baumgartner JC, Clark SM. Effects of a carbon dioxide laser on human root dentin. *J Endod* 1995;21:4-8.
56. Qadri T, Miranda L, Tunér J, Gustafsson A. The short-term effects of low-level lasers as adjunct therapy in the treatment of periodontal inflammation. *J Clin Periodontol* 2005;32:714-719.
57. Kreisler M, Meyer C, Stender E, Daubländer M, Willershausen-Zönnchen B, d'Hoedt B. Effect of diode laser irradiation on the attachment rate of periodontal

- ligament cells: An in vitro study. *J Periodontol* 2001;72:1312-1317.
58. Kreisler M, Daubländer M, Willershausen-Zönnchen B, d'Hoedt B. Effect of diode laser irradiation on the survival rate of gingival fibroblast cell cultures. *Lasers Surg Med* 2001;28:445-450.
  59. Kreisler M, Christoffers AB, Willershausen B, d'Hoedt B. Effect of low-level GaAlAs laser irradiation on the proliferation rate of human periodontal ligament fibroblasts: An in vitro study. *J Clin Periodontol* 2003;30:353-358.
  60. Ozawa Y, Shimizu N, Abiko Y. Low-energy diode laser irradiation reduced plasminogen activator activity in human periodontal ligament cells. *Lasers Surg Med* 1997;21:456-463.
  61. Heijl L. Periodontal regeneration with enamel matrix derivative in one human experimental defect. A case report. *J Clin Periodontol* 1997;24:693-696.
  62. Sculean A, Schwarz F, Chiantella GC, Arweiler NB, Becker J. Nine-year results following treatment of intrabony periodontal defects with an enamel matrix derivative: Report of 26 cases. *Int J Periodontics Restorative Dent* 2007;27:221-229.
  63. Sculean A, Schwarz F, Miliauskaitė A, et al. Treatment of intrabony defects with an enamel matrix protein derivative or bioabsorbable membrane: An 8-year follow-up split-mouth study. *J Periodontol* 2006;77:1879-1886.
  64. Silvestri M, Rasperini G, Euwe E. Enamel matrix derivative in treatment of infrabony defects. *Pract Periodontics Aesthet Dent* 1999;11:615-616, 618.
  65. Mariotti A. Efficacy of chemical root surface modifiers in the treatment of periodontal disease. A systematic review. *Ann Periodontol* 2003;8:205-226.
  66. Caffesse RG, Holden MJ, Kon S, Nasjleti CE. The effect of citric acid and fibronectin application on healing following surgical treatment of naturally occurring periodontal disease in beagle dogs. *J Clin Periodontol* 1985;12:578-590.
  67. Lowenguth RA, Blieden TM. Periodontal regeneration: Root surface demineralization. *Periodontol 2000* 1993;1:54-68.
  68. Misra V, Mehrotra KK, Dixit J, Maitra SC. Effect of a carbon dioxide laser on periodontally involved root surfaces. *J Periodontol* 1999;70:1046-1052.
  69. Schwarz F, Sculean A, Berakdar M, Szathmari L, Georg T, Becker J. In vivo and in vitro effects of an Er:YAG laser, a GaAlAs diode laser, and scaling and root planing on periodontally diseased root surfaces: A comparative histologic study. *Lasers Surg Med* 2003;32:359-366.
  70. Trylovich DJ, Cobb CM, Pippin DJ, Spencer P, Killoy WJ. The effects of the Nd:YAG laser on in vitro fibroblast attachment to endotoxin-treated root surfaces. *J Periodontol* 1992;63:626-632.
  71. Steffensen B, Webert HP. Relationship between the radiographic periodontal defect angle and healing after treatment. *J Periodontol* 1989;60:248-254.
  72. Cortellini P, Tonetti MS. Focus on intrabony defects: Guided tissue regeneration. *Periodontol 2000* 2000;22:104-132.
  73. Mangot D. The LANAP protocol: The legend continues. *Dent Today* 2013;32:130, 132.
  74. Cobb CM. Lasers in periodontics: Use and abuse. *Compend Contin Educ Dent* 1997;18: 847-852, 854-845, 858-849; quiz 860.
- Correspondence: Dr. Hom-Lay Wang, Department of Periodontics and Oral Medicine, School of Dentistry, University of Michigan, 1011 North University Ave., Ann Arbor, MI 48109-1078. Fax: 734/936-0374; e-mail: homlay@umich.edu.
- Submitted April 2, 2015; accepted for publication July 23, 2015.